arXiv:1207.5472v1 [gr-qc] 23 Jul 2012

Cosmological evolution, future singularities and Little Rip in viable $f(R)$ theories and their scalar-tensor counterpart

Diego Sáez-Gómez*
Fisika Teorikoaren eta Zientziaren Historia Saila, Zientzia eta Teknologia Fakultatea,
Euskal Herriko Unibertsitatea, 644 Posta Kutxatila, 48080 Bilbao, Spain, EU

Modified $f(R)$ gravity is one of the most promising candidates for dark energy, and even for the unification of the whole cosmological evolution, including the inflationary phase. Inside this class of theories, the so-called viable modified gravities represent realistic theories that are capable to reproduce late-time acceleration, and satisfy strong constraints at local scales, where General Relativity is recovered. In the present work, the cosmological evolution for some of these models is analyzed, which indicates that these $f(R)$ theories may lead to a phantom phase in the universe evolution. Furthermore, the scalar-tensor equivalence of $f(R)$ gravity is also considered, which can provide important properties on this kind of models. Moreover, the possibility of the occurrence of future singularities as well as the so-called *Little Rip* are studied.

PACS numbers: 04.50.Kd, 95.36.+x, 98.80.-k

I. INTRODUCTION

Since the discovery by two independent groups of the accelerating expansion of the universe through observations of the luminosity distance of Supernovae IA, as well as by some other independent observations (such as the Cosmic Microwave anisotropies), a great effort has been made to understand what kind of mechanism is producing this anomalous behavior on the expansion of the universe, encompassing the mystery under the name of dark energy. Since the value of the observed dark energy density is so large in comparison with the one predicted by quantum field theories for the vacuum energy density, it is quite reasonable to think that an unknown contribution is affecting the cosmological expansion, either by a new field such as quintessence models or vector fields, or by some kind of modification of General Relativity (GR) which has effects only on cosmological scales (for a review on modified gravities see [1]).

Then, as a consequence, along recent years the study of modified gravity theories has become very popular, and a major point for part of the scientific community, in order to understand the problem and the possibilities to reconstruct a gravitational theory capable to reproduce the dark energy epoch, and even the inflationary phase. In this sense, $f(R)$ theories seem to be the best candidate because of their simplicity, generalizing the Hilbert-Einstein action to a more complex function of the Ricci scalar, and the ease of reconstructing the corresponding action that reproduces a particular cosmological evolution (see Refs.[2–6]). Some of these models can reproduce exactly the same behavior as ΛCDM model (see Ref. [4]), or mimic a cosmological constant at the current epoch, unifying also the inflationary phase (see [5, 7]). Furthermore, as suggested by observations, the possibility that the dark energy behaves as a phantom fluid, i.e. its equation of state (EoS) is less than -1 right now or in the near future, is not excluded and has been widely explored (see Ref. [8, 9]). In such a case, the universe evolution may enter in a phantom phase, producing a super-accelerating expansion that may lead to some kind of future singularity, as the so-called *Big Rip* (for a classification of future singularities, see Ref. [10]). In this sense, $f(R)$ models that cross the phantom barrier and lead to future singularities have been also studied in Ref. [11].

Nevertheless, the high precision of local gravity tests makes that any modification of GR should pass very strong constraints, and recover GR in a certain limit (see Ref. [12]). In this sense, some viable $f(R)$ models have been proposed, which are capable to reproduce a realistic cosmological evolution, and prevent violations of local gravitational tests and instabilities in the presence of matter distributions (see Refs. [13–15]). Furthermore, some of these models are capable to reproduce both the dark energy epoch as well as the inflationary phase (see Ref. [16]). However, viable $f(R)$ gravity may lead to a phantom phase in the present/future time, which may produce some kind of future singularity (see Ref.[17]). Moreover, in the era of precise cosmology, a deeper study on the large number of dark energy models is completely mandatory in order to solve a problem of degeneracy, since most of dark energy candidates can reproduce the same cosmological evolution with no significant differences. In the case of modified gravity, some studies about the stability of the cosmological solutions [18], or the exploration of cosmological

---

* E-mail: diego.saez@ehu.es

perturbations [19], can provide some restrictions on the form of the gravitational action.

In the present work, we focus on the study of the so-called viable $f(R)$ models, exploring their cosmological evolution, and its scalar-tensor representation. It is well known that $f(R)$ gravity can be redefined as a kind of Brans-Dicke theory in terms of a scalar field with a null kinetic term (see Refs. [6, 20] and references therein), so the phase space for Friedmann-Lemaître-Robertson-Walker (FLRW) metrics can provide some useful information about the stability of the cosmological evolution as well as the presence of future singularities in viable $f(R)$ gravity. In this sense, we study two characteristic models of the class of viable $f(R)$ theories (see Refs. [13, 14]), where the cross of the phantom barrier appears as a common factor for both models, and where the possibility of a future singularity is studied. In addition, the possibility of the so-called *Little Rip* is also explored, this is a postulated phase of the universe evolution, free of singularities, where a very strong accelerating expansion may lead to break some bounded systems, as planetary systems, [21]. Such possibility may introduce some important predictions (and constraints) in the gravitational models as it may produce instabilities on the planets orbits in the near future. The *Little Rip* has been previously considered for several dark energy models [22], as well as in some contexts of modified gravities, [23]. Here we study the possibility of a *Little Rip* in viable $f(R)$ theories, in particular we study the effects on the Earth-Sun system.

The paper is organized as follows: section II is devoted to the introduction of viable $f(R)$ theories and their main cosmological properties. In section III, we explore the evolution of the effective EoS parameter of the universe, showing that in general the phantom barrier is crossed. Section IV deals with the analogous scalar-tensor theory for $f(R)$ gravity and the behavior of the phase space. Finally, in section V, we discuss the possibility of the presence of cosmological singularities according to the results found in previous sections, as well as the occurrence of a Little Rip along the cosmological evolution, where we study the effects of the expansion on the gravitational coupling in the Earth-Sun system.

II. VIABLE $f(R)$ THEORIES

Let's start by reviewing reliable cosmological models for the kind of modified gravities called viable $f(R)$ theories. We consider the action corresponding to one of these theories which, aside from the gravity part, also contains a matter contribution, namely

$$S = \int d^4x\sqrt{-g}\left[f(R) + 2\kappa^2\mathcal{L}_m\right] \ . \tag{1}$$

Here the coupling constant is given by $\kappa^2 = 8\pi G$, while $\mathcal{L}_m$ stands for the Lagrangian corresponding to the matter that fills the particular system to be studied. Note that the Hilbert-Einstein action is recovered for $f(R) = R$ in (1), The field equations corresponding to action (1) are obtained by the variation of this action with respect to the metric tensor $g_{\mu\nu}$, what yields

$$R_{\mu\nu}f_R(R) - \frac{1}{2}g_{\mu\nu}f(R) + g_{\mu\nu}\Box f_R(R) - \nabla_\mu\nabla_\nu f_R(R) = \kappa^2 T^{(m)}_{\mu\nu} \ . \tag{2}$$

Here the subscript $_R$ denotes derivatives with respect to R. We assume a flat FLRW metric,

$$ds^2 = -dt^2 + a(t)^2\sum_{i=1}^{3} dx^{i2} \ . \tag{3}$$

Then, for this metric, modified FLRW equations are obtained through the 00 and ij components of the field equations (2),

$$\begin{aligned} H^2 &= \frac{1}{3f_R}\left[\kappa^2\rho_m + \frac{Rf_R - f}{2} - 3H\dot{R}f_{RR}\right] \ , \\ -3H^2 - 2\dot{H} &= \frac{1}{f_R}\left[\kappa^2 p_m + \dot{R}^2 f_{RRR} + 2H\dot{R}f_{RR} + \ddot{R}f_{RR} + \frac{1}{2}(f - Rf_R)\right] \ , \end{aligned} \tag{4}$$

where dots denote derivatives with respect the time, Hubble parameter is defined as usual $H(t) = \dot{a}/a$, and Ricci scalar for the metric (3) is $R = 6\ (2H^2 + \dot{H})$. Here, we are interested in studying a subclass of modified gravities, the so-called viable $f(R)$ gravities, as they accomplish some indispensable conditions to be considered realistic candidates to describe the universe evolution. This kind of $f(R)$ gravities are usually described by actions of the type,

$$f(R) = R + F(R) \ . \tag{5}$$

This action basically represents Hilbert-Einstein action plus an additional term that should have only effects at cosmological scales, while at local scales, General Relativity should be recovered. In order to avoid serious problems with known physics, one has to choose the $F(R)$ function such that the theory contains flat solutions and passes also local gravity tests (see [12]). Viable gravities are able to satisfy these constraints, and to avoid large instabilities in the presence of matter distributions, as well as the anti-gravity regime by imposing positivity in the first derivative of the action, $1+F_R>0$, and the appropriate form of the function $F(R)$ (see [13]-[15]). Moreover, on the cosmological level, these theories are capable to reproduce the effects of dark energy, and even inflation. Note that both equations in (4) are written in such a way that $F(R)$-terms are put on the matter side (see Ref. [6]); thus we may define an energy density for the extra $F(R)$-terms. Hence, the first FLRW equation in (4) can be rewritten for the kind of actions (5) in terms of the cosmological parameters Ω_i,

$$1=\Omega_m+\Omega_{F(R)}\ , \quad \text{where}$$

$$\Omega_m=\frac{\rho_m}{\frac{3}{\kappa^2}H^2}\ , \quad \Omega_{F(R)}=\frac{1}{3H^2}\left(\frac{RF_R-F}{2}-3H\dot{R}F_{RR}-3H^2F_R\right)\ . \tag{6}$$

Then, the first Friedmann equation (4) takes a simple form, with two fluids contributing to the scale factor dynamics. In addition, the continuity equation $\nabla_\mu T^{\mu\nu}=0$ for a perfect fluid with an EoS $p_m=w_m\rho_m$ yields,

$$\dot{\rho}_m+3H(1+w_m)\rho_m=0\ . \tag{7}$$

Hence, for a particular $F(R)$, the corresponding cosmological evolution can be obtained through equations (4), and the continuity equation (7).

In absence of any kind of matter, dynamics of the universe are carried out by the $F(R)$-component, which may be chosen so that reproduces (or at least contributes) to the early inflationary phase and the late-time accelerating epoch. To reproduce the whole history of the universe, the following conditions on the $F(R)$ function have been proposed (see [14]):

i) Inflation occurs under one of the following conditions:

$$\lim_{R\to\infty}F(R)\to-\Lambda_i \quad \text{or} \quad \lim_{R\to\infty}F(R)\to\alpha R^n\ . \tag{8}$$

In the first situation in (8), $F(R)$ function behaves as an effective cosmological constant at early times, while the second condition yields also an accelerating expansion where the scale factor behaves as $a(t)\sim t^m$, with $m=\frac{2n^2+1-3n}{2-n}$ (see Ref. [3]).

ii) In order to reproduce late-time acceleration, we can impose a similar condition on $F(R)$. In this case, Ricci scalar has a finite value, which is assumed to be the current one, so that the condition is expressed as

$$F(R_0)=-2R_0 \quad F'(R_0)\sim 0\ . \tag{9}$$

This basically means that extra terms in the action behave as an effective cosmological constant at the present time, $p\sim-\rho$. However, as the effective fluid, coming from extra terms in the action, is not exactly a cosmological constant, the evolution would be quite different, since the universe may enter in a phantom phase [17], or may produce oscillations along the cosmological evolution [24]. We will explore the evolution, and the crossing of the phantom barrier in viable $f(R)$ gravity in next sections.

Hence, under these circumstances, the $F(R)$ term is able to reproduce both accelerating epochs of the universe evolution. Here we will focus on the study of two models of this kind of viable gravity, proposed by Hu and Sawicki in Ref. [13], and Nojiri and Odintsov in Ref. [14], whose actions are given by,

$$F_{HS}(R)=-R_{HS}\frac{c_1(R/R_{HS})^n}{c_2(R/R_{HS})^n+1}\ , \quad F_{NO}(R)=\frac{R^n(aR^n-b)}{1+cR^n}\ . \tag{10}$$

Here, $\{c_1,c_2,n\}$ are free parameters and $R_{HS}=\kappa^2\rho_m^0$ according to Ref. [13], while $\{a,b,c,n\}$ are free parameters for the second model in (10). The first model in (10) has been studied in Ref. [7], where it was proven that the universe evolution goes through two different De Sitter points, being one of them stable and the other one unstable, which can be identified to the current accelerated era and to the inflationary epoch, respectively. Similarly, the second model in (10) was studied in Ref. [16], also with the presence of an extra field, and was shown that can well reproduce both accelerated epochs, mimicking a cosmological constant at the present time. Thus, the models (10) are capable to reproduce both accelerated epochs, but a different cosmological evolution than ΛCDM model is reproduced. In next sections, we will study the evolution of both models, specially during the dark energy epoch, focusing on the possibility of the crossing of the phantom barrier. We will also study the behavior of their scalar-tensor equivalent, where the phase space will be explored.

III. COSMOLOGICAL EVOLUTION IN VIABLE $f(R)$ GRAVITY

In this section, we explore the cosmological evolution for the models considered above. For that reason, it is convenient to perform a change of variables in the equations (4), considering the redshift z as the independent variable instead of the cosmological time t, where the redshift is defined as usual,

$$1+z=\frac{a_0}{a(t)}\ , \tag{11}$$

and a_0 is the value of the scale factor at the present time t_0, such that the current epoch corresponds to $z=0$ Then, the time derivative is transformed as $\frac{d}{dt}=-(1+z)H\frac{d}{dz}$, and the first FLRW equation in (4) and the continuity equation (7) yield,

$$H^2(z)=\frac{1}{3f_R}\left[\kappa^2\rho_m(z)+\frac{R(z)f_R-f}{2}+3(1+z)H^2f_{RR}R'(z)\right]\ , \tag{12}$$

$$(1+z)\rho_m'(z)-3(1+w_m)\rho_m(z)=0\ , \tag{13}$$

where now primes denote derivatives with respect to the redshift. Then, the Ricci scalar can be rewritten as $R=6\left[2H^2(z)-(1+z)H(z)H'(z)\right]$, while the equation (13) can be easily solved for a constant EoS parameter w_m,

$$\rho(z)=\rho_0(1+z)^{3(1+w_m)}\ . \tag{14}$$

Here ρ_0 is the value of the matter energy density at the present epoch $z=0$, which can be rewritten as $\rho_0=\frac{3}{\kappa^2}H_0^2\Omega_m^0$. Then, we can fit the current values of the cosmological parameters using the observational data [26], where $H_0=100\,h$ km s^{-1} Mpc^{-1} with $h=0.71\pm0.03$ and the matter density, $\Omega_m^0=0.27\pm0.04$, while the matter fluid is considered pressureless (cold dark matter and baryons), such that $w_m=0$. The equation (12) is a second order differential equation on $H(z)$, so by fixing the initial conditions, the corresponding cosmological evolution can be obtained through the Hubble parameter in terms of the redshift for both models considered in previous section, and explore how the future evolution (with $-1<z<0$) might be. Here, we explore the evolution of the effective equation of state parameter w_{eff}, which is defined as

$$w_{eff}=\frac{p_{F(R)}+p_m}{\rho_{F(R)}+\rho_m}=-1-\frac{2\dot{H}(t)}{3H^2(t)}=-1+\frac{2(1+z)H'(z)}{3H(z)}\ . \tag{15}$$

For simplicity we redefine the Hubble parameter as $H(z)=H_0\ h(z)$, such that $h(0)=1$, while the initial condition on the first derivative $h'(0)$ can be fixed by extrapolating a particular model at $z=0$. In this sense, if we assume that ΛCDM is a good description at $z=0$, FLRW equations are given by,

$$H^2=\frac{\kappa^2}{3}\rho_m+\frac{\Lambda}{3}\ ,\quad \dot{H}=-\frac{\kappa^2}{2}\rho_m\ . \tag{16}$$

Performing the change of variables $t\to z$, and $H(z)=H_0\ h(z)$, the second equation in (16) at $z=0$ yields,

$$h'(z)=\frac{\kappa^2}{2}\frac{\rho_m}{H_0^2\ (1+z)h(z)}\ ,\quad\to\quad h'(0)=\frac{\kappa^2}{2H_0^2}\rho_m^0=\frac{3}{2}\Omega_m^0\ . \tag{17}$$

We are interested to explore the cosmological evolution for the models (10) by assuming that the universe mimics ΛCDM model (16) at present time ($z=0$), so that the initial conditions (17) are assumed. Moreover, we may compare such assumption with the one that supposes the universe has already crossed (or will do very soon) the phantom barrier, i.e. the universe is already dominated by a kind of phantom fluid at $z=0$, a possibility allowed by observations. In such a case, the first derivative of the Hubble parameter with respect to the redshift should be approximately zero or negative, $h'(0)\lesssim 0$.

Let's now analyze the $F_{HS}(R)$ model in (10), assuming $c_1=2$ and $c_2=1$, which are dimensionless parameters, and a power of $n=1$. In Fig. 1, the evolution of the EoS parameter (15) is plotted with respect to the redshift for initial conditions that mimic ΛCDM model at $z=0$. In spite of the evolution of the EoS parameter is quite similar to the ΛCDM model for positive redshifts, the universe enters in a phantom phase in the future (negative redshifts), where the EoS parameter turns out less than -1, and it oscillates close to $z=-1$ as shown in fig. 1b in more detail. In Fig. 2, the EoS parameter (15) is also evaluated, but assuming different initial conditions at $z=0$. In this case, the



(a) (b)

Figure 1: Evolution of the EoS parameter (15) as a function of the redshift z for the model $F_{HS}(R)$ where $n = 1$, $c_1 = 2$, $c_2 = 1$, and initial conditions $h(0) = 1$ and $h'(0) = 0.4$, according to (17). Figure 1a shows the evolution from the past to the future, where the phantom barrier is crossed for negative redshifts. In fig. 1b, we can see in more detail the range $-1 < z < -0.8$, where the phantom barrier is clearly crossed, and the EoS oscillates.



(a) (b)

Figure 2: Evolution of w_{eff} for the model $F_{HS}(R)$, where $n = 1$, $c_1 = 2$, $c_2 = 1$, and assuming that the universe has already entered in the phantom phase, $h(0) = 1$ and $h'(0) = -0.1$.As shown in figure 2a, the EoS crosses the phantom barrier even before $z = 0$, and it oscillates along the barrier in the future. In more detail, we can see the future evolution in fig. 2b, which ends outside of the phantom region

universe has already entered in the phantom phase at $z = 0$. The evolution for the second case, plotted in Fig. 2, shows that the amplitude of the oscillations start before, and they are larger than in Fig. 1, while the evolution for redshifts $z > 0.5$ is quite similar. Hence, both examples can give accurate descriptions of the universe evolution in the past, but they predict clearly different behaviors for the future. Furthermore, both examples present oscillations around the phantom barrier when the redshift is close to $z = -1$, as shown by Figs. 1b-2b.

In Figs. 4 and 5, the evolution of the EoS parameter for the NO model (10) is evaluated. As in HS model, firstly we assume initial conditions given by $h'(0) = \frac{3}{2}\Omega_m^0$ (Fig. 3), and then, in Fig. 4, $h'(0) \sim -0.1$. For NO model, not all the free parameters are dimensionless as above, but $\{a, c\}$ have dimensions of H^{-2} while b is effectively dimensionless.



(a) (b)

Figure 3: Evolution of the EoS parameter for $F_{NO}(R)$. We have assumed $n = 1$, $a = 0.1/H_0^2$, $b = 1$, $c = 0.05/H_0^2$ and initial conditions similar to ΛCDM model at $z = 0$, $h(0) = 1$ and $h'(0) = 0.4$. As shown, this model behaves very similarly to Hu-Sawicki model, fig. **??** is similar to fig. 1a, while the graphic 3b shows that the future evolution enters in the phantom region, and bounces above the barrier, with no oscillations.



(a) (b)

Figure 4: Evolution of $w_{eff}(z)$ for $F_{NO}(R)$, with $n = 1$, $a = 0.1/H_0^2$, $b = 1$, $c = 0.05/H_0^2$, and $h(0) = 1$ and $h'(0) = -0.1$. Figure 4a shows that the evolution oscillates and the universe crosses the phantom barrier before $z = 0$, while the oscillations remain above -1 for $-1 < z < 0$ as shown in more detail in Fig. 4b.

As shown in Fig. 3, the behavior of NO model is similar to the HS viable model: the EoS evolutes similar as ΛCDM model for $z > 0$, while it enters in the phantom region when $z < 0$ where main differences with respect the above model are found. According to the Fig. 3b, the phantom barrier is crossed, but then, it bounces above $w = -1$, and no oscillations are produced as in Fig. 1b. For the second case (Figs. 4), where the crossing of the phantom barrier is assumed to occur before $z = 0$, the EoS parameter for NO model oscillates, and the universe crosses the phantom barrier in positive redshifts, ending above the barrier $w = -1$, similarly as HS model, shown in Fig. 2a.

Figures 5 and 6 show the evolution of the cosmological parameters $\{\Omega_m, \Omega_F\}$ with respect to the redshift for the Hu-Sawicki and Nojiri-Odintsov models respectively for the different initial conditions considered above. Note that



(a)

(b)

Figure 5: Evolution of cosmological parameters $\{\Omega_m, \Omega_F\}$ defined in (6) as a function of the redshift z for the HS model with $n = 1$, $c_1 = 2$, $c_2 = 1$, and assuming initial conditions, $h(0) = 1$ and $h'(0) = 0.4$ (fig. 5a), and $h(0) = 1$ and $h'(0) = -0.1$ (fig5b). Both evolutions seem very similar, but in the right one, the cosmological parameter Ω_F decays for $0 < z < 1$, but it starts to increase again for $z > 1$, clearly in contradiction with standard model.



(a)

(b)

Figure 6: Evolution of cosmological parameters $\{\Omega_m(z), \Omega_F(z)\}$ for $F_{NO}(R)$ with $n = 1$, $a = 0.1/H_0^2$, $b = 1$, $c = 0.05/H_0^2$, and assuming initial conditions, $h(0) = 1$ and $h'(0) = 0.4$ (fig. 6a), and $h(0) = 1$ and $h'(0) = -0.1$ (fig. 6b). In fig. 6b, the transition to the dark energy epoch occurs before than in fig. 6a, a consequence of the initial conditions.

the choice of the initial conditions affect the cosmological evolution for both models, while for the HS model, both cases present the same evolution for negative redshifts, for positive ones, the evolution is different as shown in Fig. 5b, where the cosmological parameters seem to oscillate when, $z > 0$. In the NO model, when phantom initial conditions are assumed, Ω_F tends to dominate completely the universe, while for positive redshifts, it becomes negative, fig. 6b.

Note that, as shown in figures 1-4, viable modified gravities (10) produce some oscillations along the cosmological evolution, a fact pointed out before for this kind of models in Ref. [24]. In general, we have shown that the transition to the phantom epoch occurs. Moreover, the election of phantom conditions at $z = 0$ gives an anomalous behavior of the cosmological parameters for both models when $z > 0$.

IV. SCALAR-TENSOR REPRESENTATION OF $f(R)$ GRAVITY

It is well known that $f(R)$ gravity is equivalent to a kind of Brans-Dicke theory with a null kinetic term, and a scalar potential, (see for example, [6, 20] and references therein),

$$S = \int d^4x\sqrt{-g}\left[\phi\, R - V(\phi) + 2\kappa^2\mathcal{L}_m\right]\ . \tag{18}$$

By varying the action with respect to the scalar field ϕ, the relation between both theories is obtained, and $f(R)$ action (1) is recovered,

$$R = V'(\phi) \quad \rightarrow \quad \phi = \phi(R)\ ,\ \Rightarrow \quad f(R) = \phi(R)R - V(\phi(R))\ . \tag{19}$$

While the scalar field and the potential are related with the particular $f(R)$ action by,

$$\phi = f_R(R)\ ,\quad V(\phi(R)) = f_R(R)\ R - f(R)\ . \tag{20}$$

Then, by a particular $f(R)$ action, we can reconstruct its equivalent scalar-tensor theory (18), so that we can study the viable models studied above by analyzing their scalar-tensor counterpart. Moreover, by varying the action (18) with respect to the metric $g_{\mu\nu}$, field equations are obtained,

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{\kappa^2}{\phi}T^{(m)}_{\mu\nu} + \frac{1}{\phi}(\nabla_\mu\nabla_\nu\phi - g_{\mu\nu}\Box\phi) - \frac{1}{2}g_{\mu\nu}V(\phi)\ ,$$

$$3\Box\phi = \kappa^2T^{(m)} + \phi\frac{dV(\phi)}{d\phi} - 2V(\phi)\ , \tag{21}$$

where the second equation is the trace of the field equations in (21). Hence, for a flat FLRW metric (3), the Friedmann equations yield,

$$3H^2 = \frac{1}{\phi}\left(\kappa^2\rho_m - 3H\dot{\phi} + \frac{1}{2}V(\phi)\right)\ ,\quad -3H^2 - 2\dot{H} = \frac{1}{\phi}\left(\kappa^2p_m + \ddot{\phi} + 2H\dot{\phi} - \frac{1}{2}V(\phi)\right)\ . \tag{22}$$

while the trace equation in (21) is given by,

$$3\ddot{\phi} = \kappa^2(\rho_m - p_m) - \phi V' + 2V - 9H\dot{\phi}\ . \tag{23}$$

Let's reconstruct the corresponding scalar-tensor theory for the particular models considered in the previous section. By the expressions (20), the corresponding relation $\phi(R)$ for the models (10) are given by,

$$\phi_{HS} = 1 + \frac{c_1c_2R}{R_{HS}\left(1+\frac{c_2R}{R_{HS}}\right)^2} + \frac{c_1}{1+\frac{c_2R}{R_{HS}}}\ ,\quad \phi_{NO} = 1 - \frac{cR(aR-b)}{(1+cR)^2} + \frac{aR}{1+cR} + \frac{aR-b}{1+cR}\ , \tag{24}$$

where we have assumed a power of $n = 1$ for both models (10). Then, the scalar potentials (20) yield,

$$V_{HS}(\phi) = \frac{1+c_1-\phi\pm 2\ c_2\sqrt{c_1\ (1-\phi)}}{c_2}R_{HS}\ ,\quad V_{NO}(\phi) = \frac{2\ a + c\ (1+b-\phi)\pm 2\ \sqrt{(a+b\ c)\ (a+c-\phi)}}{c^2}\ . \tag{25}$$

Hence, the scalar-tensor representation for both models is not uniquely defined, but the potentials exhibit two branches, which in principle do not affect the cosmological evolution, but it may influence the behavior of the phase space. In addition, both models introduce a boundary condition on the value of the scalar field, being $\phi < 1$ for the HS model, and $\phi < a + c/c$ for the NO model, a limit where both branches of the potentials converge, as can be seen in Fig. 7.

Moreover, the evolution of the scalar field as a function of the redshift, $\phi(z) = f_R(R(z))$, is the same regardless of the sign in the potentials (25), as shown in Fig. 8. Note that there is a direct correlation between the behavior of the scalar field and the evolution of the EoS parameter as can be shown comparing figs. 1 and 3 with fig. 8. For the HS model, the EoS parameter (Fig. 1)presents some oscillations, specially for negative redshifts around the phantom barrier, while the evolution of the scalar field also oscillates according to Fig. 8a. In the NO model, the linear behavior of the EoS parameter observed in Fig. 3 is also *translated* in a linear evolution of the scalar field with respect to the redshift, see fig. 8b. Let us now analyze the phase space $\{H, \phi\}$ for this kind of viable models in vacuum, where



(a) (b)

Figure 7: Scalar potentials $V(\phi)$ in terms of H_0^2, of the HS model with $n = 1$, $c_1 = 2$, $c_2 = 1$ (figure 7a), and of the NO model with $n = 1$, $a = 0.1/H_0^2$, $b = 1$, $c = 0.05/H_0^2$, fig. 7b. Both potentials are not uniquely defined, where each branch behaves very different, converging to the boundary of the scalar field ϕ.



(a) (b)

Figure 8: Evolution of the scalar field $\phi(z)$ for the HS model with $n = 1$, $c_1 = 2$, $c_2 = 1$ (a), and for the NO model (b) with $n = 1$, $a = 0.1/H_0^2$, $b = 1$, $c = 0.05/H_0^2$. In both cases, the initial conditions are: $h(0) = 1$, and $h'(0) = 3\Omega_m^0/2$. For each example, we can easily see the correlation between the evolution of the EoS parameter in Figs. 1 and 3, and the evolution of the scalar field for each model respectively

both branches of the scalar potentials are considered. For simplicity, we assume vacuum FLRW equations, so that by combining equations (22)-(23), we get,

$$\dot{H} = -2H^2 + \frac{1}{6}V'(\phi) \ , \quad \dot{\phi} = \frac{1}{3H}\left[-3H^2\phi + \frac{1}{2}V(\phi)\right] \ . \tag{26}$$

Hence, by combining both equations in (26),

$$\frac{dH}{d\phi} = \frac{\frac{V'(\phi)}{6} - 2H^2}{\frac{V(\phi)}{3H} - H\phi} \ . \tag{27}$$



(a) (b)

Figure 9: Phase space $H(\phi)$ for the HS model, where different initial conditions are considered, and both branches of the potential are studied. The left figure corresponds to $V_+(\phi)$ and the right one to $V_-(\phi)$. In both cases, we have considered initial conditions close to the boundary of the scalar field: (a) $H(0.95) = H_0^2$, (b) $H(0.95) = 10H_0^2$, (c) $H(0.95) = 100H_0^2$.



(a) (b)

Figure 10: Phase space $H(\phi)$ for the NO model. As above, different initial conditions are considered, and both branches of the potential are studied. The left figure corresponds to $V_+(\phi)$ and the right one to $V_-(\phi)$. We have considered initial conditions close to the boundary of the scalar field: (a) $H(2.95) = H_0^2$, (b) $H(2.95) = 10H_0^2$, (c) $H(2.95) = 100H_0^2$.

This equation describes the phase space for a particular scalar potential, and provide the information about the behavior and stability of a particular model. Specifically, we can see in Fig. 9 that regardless of the choice of initial conditions, the behavior of the Hubble parameter tends to a stable point, close to zero. However, as shown above, the scalar field takes values very distant of the stable point, so the sensibility to the initial conditions can be important along the cosmological evolution naturally, although the evolution presents the same tendency independently of the initial conditions. In fact, we can observe some small fluctuations along the evolution of H with respect to ϕ. Moreoever, both branches of the potential V_{HS} in (25) may present different convergence points in spite of the differences may not be detected in the range of values that the scalar field takes along the cosmological history(see Fig. 8). For the NO model in Fig. 10, a similar behavior is found, where different convergence points are found for $V_\pm$, as well as larger fluctuations for V_+, while the behavior is similar in both cases, and independently on the initial

conditions (assumed close to the boundary of the scalar field).
Hence, the cosmological evolution in viable $f(R)$ gravity shows a similar behavior regardless of the model, and even of the initial conditions, where the crossing of the phantom barrier is a common element in all of them, but with no sign of possible future singularities, such that viable $f(R)$ gravity represents a serious candidate for explaining the dark energy epoch. The tiny differences between models seem very clear when the future evolution of the EoS is studied, and whose phase space does not seem to provide useful information for its discrimination. However, the stability shown in the phase space for both models suggest that no future singularity will occur in the future. In next section, we discuss this point in more detail, and we analyze the possibility of the occurrence of a *Little Rip*.

V. FUTURE SINGULARITIES AND LITTLE RIP IN VIABLE $f(R)$ GRAVITY

In recent years, the study of future singularities has become a major task since a lot of cosmological models, capable to describe a realistic cosmological evolution satisfying observational constraints, lead to the so-called phantom phase, $w < -1$, and may end in some type of future singularity (see Refs. [8]). A classification of future singularities was presented in Ref. [10],

- Type I ("Big Rip"): For $t \to t_s$, $a \to \infty$ and $\rho \to \infty$, $|p| \to \infty$.
- Type II ("Sudden"): For $t \to t_s$, $a \to a_s$ and $\rho \to \rho_s$, $|p| \to \infty$.
- Type III: For $t \to t_s$, $a \to a_s$ and $\rho \to \infty$, $|p| \to \infty$.
- Type IV: For $t \to t_s$, $a \to a_s$ and $\rho \to \rho_s$, $p \to p_s$ but higher derivatives of Hubble parameter diverge.

Future singularities have been widely studied, in particular, they may occur for some $f(R)$ models (see Refs. [6, 11]). Nevertheless, the fact that the universe may cross the phantom barrier at $w = -1$ is not sufficient condition to present a future singularity (see Ref. [9]). In section III, we have shown that in general, viable models of $f(R)$ gravity leads to a phantom epoch in the future, where the EoS parameter becomes less than -1, but no future singularity occurs since Hubble parameter and its first derivative remain finite, or in other words, according to the classification, there will not be a Big Rip, Sudden singularity or Type III singularity for the viable models studied above. In addition, type IV singularity is not completely excluded, but this kind of singularity, that affects to higher derivatives of the Hubble parameter, may not have any physical consequence, at least observationally as pointed out in Ref. [10], so its importance in comparison with the others is lower.

Nevertheless, the absence of future singularities for a particular model is not enough to avoid some critical consequences as the so-called *Little Rip*. The *Little Rip* is a postulated phase of the universe evolution Ref. [21], when a very strong accelerating expansion would lead to break some bounded systems, as the Solar System or the galaxies, but where all physical magnitudes remain finite (see also Ref. [22]). This special scenario has been also explored in modified gravity, where the occurrence of this scenario seems very possible for some particular models (see Refs. [23]). Here we are interested to study the possibility of a Little Rip in viable $f(R)$ theories, in particular in the models studied in previous sections. Firstly, we should characterized in some way the strength of the force of the cosmological expansion in order to compare with those forces of a bounded system, as for example the newtonian force in Solar System. In this sense, we may write the force that a body of mass m may feel due to the cosmological expansion by the relative acceleration between two points separated a comoving distance r,

$$F_{cosm} = m\ r\ \left(\frac{\ddot{a}}{a}\right) = m\ r\left(H^2 + \dot{H}\right)\ . \tag{28}$$

Then, we may compare this force with the bounding force of a particular system. For example, in the case of the Solar system, the force that maintains the coupling between Sun and a body of mass m can be characterized by the Newtonian law,

$$F_N = G\frac{mM_\odot}{r^2}\ . \tag{29}$$

where $M_\odot = 1.9\ 10^{30}\ kg$ is the mass of the Sun, and r is the distance between the Sun and the body m. Here, in both cases it is assumed that $F_i = |\vec{F}_i|$, as both forces have radial directions, but opposite sign. Hence, we may compare both quantities along the cosmological evolution,

$$\frac{F_{cosm}}{F_N} = \frac{r^3}{G\ M_\odot}H_0^2\left[h(z)^2 - (1+z)h(z)\ h'(z)\right]\ , \tag{30}$$

where recall that we defined $H(z) = H_0\, h(z)$ in section III, where $H_0 = 100\,h$ km s^{-1} Mpc^{-1} with $h = 0.71 \pm 0.03$. Let's compare both forces for the Earth-Sun system, where $r = 149.6 \times 10^9\ m$ is the Earth-Sun distance, at the present time, $z = 0$,

$$\frac{F_{cosm}}{F_N} = 1.4 \times 10^{-22} \left[h(0)^2 - h(0)h'(0) \right] \ , \tag{31}$$

Then, by assuming the initial conditions from section III, $h(0) = 1$ and $a)$ $h'(0) = \frac{3}{2}\Omega_m^{(0)}$, and $b)$ $h'(0) = -0.1$,

$$(i)\ \frac{F_{cosm}}{F_N} \sim 0.82 \times 10^{-22} \ , \quad (ii)\ \frac{F_{cosm}}{F_N} \sim 1.95 \times 10^{-22} \ . \tag{32}$$

Hence, both values are too small to produce any kind of instability on the orbits of the planets, and in particular on the Earth orbit. In addition, the future cosmological evolution does not seem to affect significantly to the Earth-Sun system whatever the $f(R)$ model is used, and independently on the initial conditions imposed at $z = 0$ as shown in Fig. 11 (Hu-Sawicki model) and Fig. 12 (Nojiri-Odintsov model). Nevertheless, some tiny differences are observed between both models. For the HS model, the ratio (30) behaves as a damped oscillator, being the amplitude reduced in the future (negative redshifts), a non expected behavior according to the evolution of the EoS parameter shown in section III. However, the amplitude is too small to affect the Newtonian force between the Earth and Sun. Moreover, assuming that the universe has already entered the phantom phase at $z = 0$ affects directly to the evolution of the ratio (30), where the force of the expansion is larger near $z = 0$, and even negative in the past, where it is assumed that the dust matter dominates. In the case of the NO model, different initial conditions affect directly to the evolution of the expansion force, being maximum in both cases near $z = 0$ and decreasing when the universe approximates to $z = -1$. In both cases of the NO model, the force becomes negative around $z = 1$, where dust matter dominates, but oscillates for small redshifts, where the amplitude is larger in the case of initial phantom conditions, fig. 12b.



(a)

(b)

Figure 11: Evolution of the ratio $\frac{F_{cosm}}{F_N}$ for the Earth-Sun system in the HS model, where $h(0) = 1$ and: (a) $h'(0) = \frac{3}{2}\Omega_m^{(0)}$, (b) $h'(0) = -0.1$.

Hence, viable $f(R)$ gravity does not seem to affect the bounding systems, as Solar System. Since the expansion force is proportional to the distance between two points, it may affect to some objects far away from the center of galaxies as are weakly coupled to them. However, this tiny effect does not seem to be really detectable in comparison to other effects.

VI. CONCLUSIONS

Along the present paper, we have studied the so-called viable $f(R)$ models by analyzing two characteristic examples. We found that the cosmological evolution of the effective EoS for the universe enters a phantom stage



(a)

(b)

Figure 12: Evolution of the ratio $\frac{F_{cosm}}{F_N}$ for the Earth-Sun system in the NO model, where $h(0) = 1$ and: (a) $h'(0) = \frac{3}{2}\Omega_m^{(0)}$, (b) $h'(0) = -0.1$..

in the present or near future, independently of the initial conditions imposed. In this analysis, we have seen that both models can reproduce quite well the cosmological evolution, similarly to ΛCDM model at $z = 0$, where we have imposed some particular conditions in order to fit the model with the observational constraints. As shown in section III, when the phantom transition is assumed to occur before $z = 0$, the behavior of the cosmological parameters $\{\Omega_m, \Omega_F\}$ when $z > 0$ present some inconsistencies for both models, such that those conditions may be excluded.
On the other hand, the transition to the phantom phase does not imply directly the occurrence of a future singularity, since the Hubble parameter and its first derivatives remain finite. The possibility of a IV type singularity is not completely excluded as it implies an infinite number of higher derivatives of the Hubble parameter, but its possible physical effects leave unaffected the cosmological evolution. Even in the case that the transition to the phantom era has already occurred at $z = 0$, future evolution seems to be free of singularities. However, the choice of the initial conditions at $z = 0$ may affect the past evolution, as shown in the right fig. 5b for the HS model, where the evolution of the cosmological parameters Ω_i seems to deviate from the correct evolution for positive redshifts.

Furthermore, by using the equivalent picture of $f(R)$ gravity as a scalar-tensor theory, we have analyzed the evolution of the scalar field in Sect. IV, we note a correlation between the cosmological evolution shown in section III, and the evolution of the scalar field. In addition, we have seen that in general, this kind of viable modified gravities are degenerated in their scalar-tensor representation, giving two potentials that impose a bound on the scalar field and tends to the same value. Nevertheless, by exploring the phase space in vacuum, we found that independently on the initial conditions, both branches tend to stable points, encouraging the idea that no future singularity occurs.

In the last section, we discussed the possibility of the occurrence of a future singularity, but also the so-called *Little Rip*, an stage of the universe evolution that may affect some bounded systems due to the strength of the expansion, but where all the physical magnitudes remain regular. We have explore this possibility by comparing the strength felt by a body with respect to another point of the universe, and the newtonian force of a typical planet system. For the Earth-Sun system, the expansion force is too small to produce any effect on the orbit. However, as pointed above, some systems weakly coupled may have some tiny effects, although the possibility to detect and distinguish those effects from others seems not possible.

Hence, at the present work, we have performed a deep analysis on viable $f(R)$ gravities, showing that the models considered here are capable to reproduce the late-time acceleration, entering in a phantom phase in the present or near future, a fact that may be a way to distinguish from other dark energy models. Furthermore, the absence of future singularities, and instabilities in the orbits of the planets, provide a strong support to the so-called viable modified gravity as a serious candidate for explaining the dark energy paradigm.

Acknowledgments

I would like to thank Sergei Odintsov, and Alvaro de la Cruz-Dombriz for useful discussions on the topic. I acknowledge support from a postdoctoral contract from the University of Basque Country (UPV/EHU). I am also supported by the research project FIS2010-15492, and also by the Basque Government through the special research action KATEA and UPV/EHU under program UFI 11/55.

---

[1] S. Nojiri and S. D. Odintsov, eConf **C0602061**, 06 (2006) [Int. J. Geom. Meth. Mod. Phys. **4**, 115 (2007)]; hep-th/0601213; arXiv: 0807.0685; K. Bamba, S. Capozziello, S. Nojiri and S. D. Odintsov, arXiv:1205.3421 [gr-qc]; S. Capozziello and M. De Laurentis, Phys. Rept. **509**, 167 (2011) [arXiv:1108.6266 [gr-qc]]; S. Capozziello and M. Francaviglia, Gen. Rel. Grav. **40**, 357 (2008) arXiv:0706.1146 [astro-ph]; V. Faraoni arXiv:0810.2602v1 [gr-qc]; F. S. N. Lobo, arXiv: 0807.1640 [gr-qc]; S. Capozziello and V. Faraoni, *Beyond Einstein Gravity*, Fundamental Theories of Physics Vol. 170, Springer Ed., Dordrecht (2011); A. de la Cruz-Dombriz and D. Sáez-Gómez, arXiv:1207.2663 [gr-qc].

[2] A. A. Starobinsky *Phys. Lett.* **91B** 99 (1980); S. Nojiri and S. Odintsov *Phys. Rev.* **D 68** 123512 (2003); *Gen. Rel. Grav.* **36** 1765 (2004); Phys. Rev. D **74**, 086005 (2006), hep-th/0608008; J. Phys. A **40**, 6725 (2007), hep-th/0610164; S. Capozziello, S. Nojiri, S. D. Odintsov and A. Troisi, Phys. Lett. B **639**, 135 (2006), astro-ph/0604431; S. Capozziello *Int. J. Mod. Phys.* **D 11**, 483 (2002); S. M. Carroll *et al*, *Phys. Rev.* **D71** 063513 (2005); S. Carloni, P. K. S. Dunsby, S. Capozziello and A. Troisi *Class. Quant. Grav.* **22** 4839 (2005); T. P. Sotiriou and S. Liberati *Annals of Physics* 332 (2007); J. A. R. Cembranos *Phys. Rev.* **D 73** 064029 (2006); T. Clifton and J. D. Barrow *Phys. Rev.* **D 72** 103005 (2005); S. Capozziello, S. Carloni and A. Troisi, Recent Res. Dev. Astron. Astrophys. **1**, 625 (2003) astro-ph/0303041; J. A. R. Cembranos, Phys. Rev. Lett. **102** 141301 (2009), arXiv:0809.1653 [hep-ph]; P. K. S. Dunsby, E. Elizalde, R. Goswami, S. Odintsov and D. Sáez-Gómez, Phys. Rev. D **82**, 023519 (2010), arXiv:1005.2205 [gr-qc]; K. Bamba, C. Q. Geng and C. C. Lee, JCAP **1008**, 021 (2010), arXiv:1005.4574 [astro-ph.CO]; L. G. Jaime, L. Patino and M. Salgado, arXiv:1206.1642 [gr-qc];

[3] N. Goheer, J. Larena and P. K. S. Dunsby, Phys. Rev. D **80**, 061301 (2009), arXiv:0906.3860 [gr-qc].

[4] A. de la Cruz-Dombriz and A. Dobado, Phys. Rev. D **74**, 087501 (2006), gr-qc/0607118.

[5] S. Nojiri, S. D. Odintsov and D. Sáez-Gómez, Phys. Lett. B **681**, 74 (2009), arXiv:0908.1269 [hep-th].

[6] D. Sáez-Gómez, Gen. Rel. Grav. **41**, 1527 (2009) arXiv:0809.1311 [hep-th]; S. Capozziello, S. Nojiri, and S.D. Odintsov, Phys. Lett. B **639** 135 (2006) hep-th/0512118.

[7] G. Cognola, E. Elizalde, S. D. Odintsov, P. Tretyakov, and S. Zerbini, Phys. Rev. D **79** 044001 (2009) arxiv:0810.4989[gr-qc].

[8] R. R. Caldwell, Phys. Lett. B **545** 23 (2002), astro-ph/9908168; R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. **91** 071301 (2003), astro-ph/0302506; B. McInnes, JHEP **0208** 029 (2002), hep-th/0112066; S. Nojiri and S. D. Odintsov, Phys. Lett. B **562** 147 (2003), arXiv:hep-th/0303117; V. Faraoni, Int. J. Mod. Phys. D 11, **471** (2002), astro-ph/0110067; P. F. Gonzalez-Diaz, Phys. Lett. B **586** 1 (2004), astro-ph/0312579; M. Sami and A. Toporensky, Mod. Phys. Lett. A **19** 1509 (2004), gr-qc/0312009; H. Stefancic, Phys. Lett. B **586** 5 (2004), astro-ph/0310904; L. P. Chimento and R. Lazkoz, Phys. Rev. Lett. **91** 211301 (2003), gr-qc/0307111; S. Capozziello, S. Nojiri and S. D. Odintsov, Phys. Lett. B **632** 597 (2006), hep-th/0507182; E. Elizalde, S. Nojiri, S. D. Odintsov, D. Sáez-Gómez and V. Faraoni, Phys. Rev. D **77** 106005 (2008), arxiv:0803.1311 [hep-th]]; Yuri Shtanov, Varun Sahni, Class.Quant.Grav. 19 (2002) L101-L107, gr-qc/0204040; D. Sáez-Gómez, Phys. Rev. D **85**, 023009 (2012), arXiv:1110.6033 [hep-th].

[9] V. Sahni and Y. Shtanov, JCAP **0311**, 014 (2003), astro-ph/0202346.

[10] S. Nojiri, S. D. Odintsov and S. Tsujikawa, Phys. Rev. D **71**, 063004 (2005) [hep-th/0501025].

[11] S. Nojiri and S. D. Odintsov, Phys. Rev. D **78**, 046006 (2008), arXiv:0804.3519 [hep-th]; K. Bamba, C. Q. Geng, S. Nojiri and S. D. Odintsov, Mod. Phys. Lett. A **25**, 900 (2010) arXiv:1003.0769 [hep-th].

[12] G.J. Olmo, Phys. Rev. D **75** 023511 (2007), gr-qc/0612047; S. Nojiri and S.D. Odintsov, Phys. Lett. B **652** 343 (2007), arXiv:0706.1378[hep-th]; Phys. Lett. B **657** 238-245 (2007), arXiv:hep-th/0707.1941; S. Capozziello, A. Stabile and A. Troisi, Phys. Rev. D **76**, 104019 (2007) arXiv:0708.0723 [gr-qc].

[13] W. Hu and I. Sawicki, Phys. Rev. D **76** 064004 (2007), arXiv:0705.1158[astro-ph].

[14] S. Nojiri and S.D. Odintsov, Phys. Rev. D **77** 026007 (2008), arXiv:0710.1738[hep-th].

[15] S.A. Appleby and R.A. Battye, arXiv:0705.3199[astro-ph]; L. Pogosian and A. Silvestri, Phys. Rev. D **77** 023503 (2008), arXiv:0709.0296[astro-ph]; S. Tsujikawa, Phys. Rev. D **77** 023507 (2008), arXiv:0709.1391[astro-ph]; A. Starobinsky, JETP Lett. **86** 157 (2007), arXiv:0706.2041 [astro-ph]; S. Capozziello and S. Tsujikawa, Phys. Rev. D **77** 107501 (2008) arXiv:0712.2268[gr-qc];

[16] E. Elizalde and D. Sáez-Gómez, Phys. Rev. D **80** 044030 (2009), arXiv:0903.2732.

[17] K. Bamba, C. Q. Geng and C. C. Lee, Int. J. Mod. Phys. D **20**, 1339 (2011) arXiv:1108.2557 [gr-qc].

[18] A. de la Cruz-Dombriz and D. Sáez-Gómez, arXiv:1112.4481 [gr-qc].

[19] S. M. Carroll, I. Sawicki, A. Silvestri and M. Trodden, New J. Phys. **8**, 323 (2006), astro-ph/0607458; Y. -S. Song, W. Hu and I. Sawicki, Phys. Rev. D **75**, 044004 (2007) astro-ph/0610532. R. Bean, D. Bernat, L. Pogosian, A. Silvestri and M. Trodden, Phys. Rev. D **75**, 064020 (2007) astro-ph/0611321; S. Carloni, P. K. S. Dunsby and A. Troisi, Phys. Rev. D **77**, 024024 (2008) arXiv:0707.0106 [gr-qc]; A. de la Cruz-Dombriz, A. Dobado and A. L. Maroto, Phys. Rev. D **77**, 123515 (2008) arXiv:0802.2999 [astro-ph]. A. Abebe, M. Abdelwahab, A. de la Cruz-Dombriz and P. K. S. Dunsby, Class. Quant.

Grav. **29**, 135011 (2012) arXiv:1110.1191 [gr-qc].
[20] T. Kobayashi and K. Maeda, Phys. Rev. D **79**, 024009 (2009) arXiv:0810.5664 [astro-ph].
[21] P. H. Frampton, K. J. Ludwick and R. J. Scherrer, Phys. Rev. D **84**, 063003 (2011) arXiv:1106.4996 [astro-ph.CO];
[22] P. H. Frampton, K. J. Ludwick, S. Nojiri, S. D. Odintsov and R. J. Scherrer, Phys. Lett. B **708**, 204 (2012) arXiv:1108.0067 [hep-th]. P. H. Frampton, K. J. Ludwick and R. J. Scherrer, Phys. Rev. D **85**, 083001 (2012) arXiv:1112.2964 [astro-ph.CO]; H. Wei, L. -F. Wang and X. J. Guo, arXiv:1207.2898 [gr-qc].
[23] S. Nojiri, S. D. Odintsov and D. Sáez-Gómez, AIP Conf. Proc. **1458**, 207 (2011), arXiv:1108.0767 [hep-th]; A. J. López-Revelles, R. Myrzakulov and D. Sáez-Gómez, Phys. Rev. D **85**, 103521 (2012) arXiv:1201.5647 [gr-qc].
[24] E. Elizalde, S. D. Odintsov, L. Sebastiani and S. Zerbini, Eur. Phys. J. C **72**, 1843 (2012) arXiv:1108.6184 [gr-qc]. K. Bamba, A. López-Revelles, R. Myrzakulov, S. D. Odintsov and L. Sebastiani, arXiv:1207.1009 [gr-qc].
[25] A. J. López-Revelles and E. Elizalde, Gen. Rel. Grav. **44**, 751 (2012) arXiv:1104.1123 [hep-th].
[26] WMAP Collaboration, D.N. Spergel, *et al* Astrophys. J. Suppl. **148** 175 (2003). E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **180**, 330 (2009) arXiv:0803.0547 [astro-ph]; E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192**, 18 (2011) arXiv:1001.4538 [astro-ph.CO];